

HERE PUT THE TITLE if you need small letters

A. A. AUTHOR¹ and A.A. AUTHOR²

¹*First address*

E-mail name1@ac.cd.ef.uk

²*Second address*

E-mail name2@ac.cd.ef.es

Abstract. Here put abstract

1. SECTION

text, The citation should be as:


Then emission of three regions can describe broad-lines: 1. BLR (some of authors for this region use therminology Very Broad Line Region – VBLR, see e.g. Brotherton et al. 1994, Corbin 1995, 1997); 2. Intermediate Line Region (ILR) and 3. NLR. Observations of nearby galaxies also show that the structure of NLR is more complex (see e.g. Veilleux 1991abc, Arribas et al. 1996).

1. 1. SUBSECTION

text

t	$\theta(^{\circ})$	$\rho('')$
2002.0	174.0	0.259
2003.0	174.7	0.262
2004.0	175.3	0.265
2005.0	175.9	0.268
2006.0	176.6	0.270

Table 1: Calculated positions for the next five years

Figure 1: The observed FeK_{α} line and its fit

$$G(x) = \int_0^{\infty} e^{-\frac{x^2}{2}} dx \quad (1)$$

References

- Hewitt, A., Burbidge, G. : 1989, *Astrophys. J. Suppl. Series*, **75**, 297.
 Mediavilla, E., Insertis, F. M. : 1989, *Astron. Astrophys.* **214**, 79.
 Netzer, H. : 1990, *Active Galactic Nuclei*, eds. R. D. Blandford, H. Netzer & L. Woltjer,
 Saas-Fee Advanced Course 20, Berlin: Springer – Verlag.
 Osterbrock, D. E. : 1989, *Astrophysics of Gaseous Nebulae and Active Galactic Nuclei*, Mill
 Valley, California.